



ADVERSE DRUG REACTIONS, INCLUDING SJS, TEN

FLUCONASOL ASSOCIATED TOXIDERMIA BULLOUS

E.I. Dukova⁽¹⁾

Medsic Clinic, Dermatology Department, Moscow⁽¹⁾

Other hand, drug toxidermia are secondary mucocutaneous complications due to ingestion or injection of a drug systemically use among the most common adverse drug reactions with around 20% of notifications drug events.

We have female patient 34 years old taking Fluconazol, prescribing by gynecologist. She had erythematous and bullous rash started after 5 minutes after taking drug. During 7 days she was treated by her gynecologist with antihistamins and steroid cream for external use. There was no new rash. Treatment without significant reduction.

At first visit she had erythematous rash with exfoliation on her axillas and bullous rash on her hand. Mucous membrane was clean, no fever. This patient has previously drug reaction for Fluconazole, that was not reported by her gynecologist.

We add dexamethasone 8-4-4-2 mg for 4 days with Chloropiramin 20 mg, and saline solution till 200mg, locally metilene blau. We got significant improvement after 4 days therapy.

Conclusion: We always remember about allergy for antibiotics and not always ready for drug reaction for Fluconazol. Clear allergy history of the patients allergy always help to prevent drug reaction.

