



ADVERSE DRUG REACTIONS, INCLUDING SJS, TEN

## ZONED POS-CHICKEN POX INDUCED BY CORTICOID AT A 9-YEAR-OLD CHILD

*K Yamoussa<sup>(1)</sup> - G Mamadou<sup>(2)</sup> - B Sirithio<sup>(1)</sup> - T Hawa<sup>(1)</sup> - D Adama<sup>(1)</sup> - F Ousmane<sup>(1)</sup>*

*Centre National D'appui À La Lutte Contre La Maladie, Cnam, Faculty Of Medecine Of Bamako, Dermatologie, Bamako<sup>(1)</sup> - Centre National D'appui À La Lutte Contre La Maladie, Cnam, Faculty Of Medecine Of Bamako, Dermatologie, Bamako<sup>(2)</sup>*

The Chicken pox and Zoned him are infectious, generally mild diseases caused by the virus zoned chicken pox (vzv), Zoned him(it) represent the recurrence located(localized) as a result of a modification of its virulence or disturbance of the immune defense system of the host. Objective: to highlight the imputability of corticoids in the arisen of bummed around in the immediate consequences(suites) of a chicken pox. Observation: it was about a child of 9ans without particular histories who consulted at first for an eruption feverish vésiculeuse ending in the diagnosis of chicken pox. Secondly for a new skin rash after an episode of Nasopharyngitis handled by the corticoid(Célestène drop and injectable Solumedrol) and antibiotic (amoxicilline acid clavulanique) allowing the diagnosis of bummed around led by corticoids. Conclusion: have always the notion of medicinal grip immunosuppresseurs especially corticoids at the children because the risk of arisen of bummed around.

Key words: zoned, post-chicken pox, corticoid, child

