



DERMATOPATHOLOGY

DERMATOPATHIC LYMPHADENITIS ASSOCIATED WITH VITILIGO-LIKE LESIONS IN HISPANIC PATIENT

*F Magri⁽¹⁾ - C Chello⁽¹⁾ - A Tammaro⁽¹⁾ - D Giordano⁽¹⁾ - Fr Parisella⁽²⁾ - F Persechino⁽³⁾ -
G De Marco⁽¹⁾ - S Persechino⁽¹⁾*

*Sant'andrea Hospital Faculty Of Medicine And Psychology University Of Rome
"sapienza", Italy, Nesmos Department, Dermatology Unit, Rome, Italy⁽¹⁾ - University Of
Queensland, Brisbane, Australia, Medical School, Brisbane, Australia⁽²⁾ - Sapienza
University Of Rome, Rome, Italy, Department Of Experimental Medicine, Rome, Italy⁽³⁾*

Background: Dermatopathic lymphadenitis, or lipomelanotic reticulosis (Pautrier-Woringer syndrome) is a benign hyperplasia of lymph nodes, histologically characterized by histiocytes proliferation and melanin deposits, which is often associated with eczematous manifestations, such as psoriasis and erythroderma.

Observation: Authors present a case of a male patient presenting with vitiligo-like lesions, idiopathic hypereosinophilia and diffuse lymphadenomegaly which was histologically reported as dermatopathic lymphadenopathy.

Key Message: This entity, which must be necessarily confirmed by histologic reports, can be associated with different types of cutaneous manifestation. It needs to be differentiated from pathologic lymphocytes proliferation, especially in adult patients.

