



A new ERA for global Dermatology 10 - 15 JUNE 2019 MILAN, ITALY

SKIN MANIFESTATIONS OF INTERNAL DISEASE

OROFACIAL GRANULOMATOSIS WITH CAVERNOUS HEMANGIOMA OF BRAIN

Jongsoo Hong $^{(1)}$ - Jong Heon Jeong $^{(1)}$ - Ji Young Lee $^{(1)}$ - Seung Ju Yun $^{(1)}$ - Seung Ho Lee $^{(1)}$ - Ai-young Lee $^{(1)}$

Dongguk University Ilsan Hospital, Dermatology, Goyang, Republic Of Korea (1)

Keyword: Cavernous hemangioma, Orofacial granulomatosis

44-year-old woman visited our clinic with repeated labial and perilabial swelling for months. She had been treated with oral antiviral agents, but not effective. Recently, she was diagnosed with a brain cavernous hemangioma. Histologic examination revealed exocytosis of some lymphocytes and dense perivascular, perifollicular lymphohistiocytic infiltration throughout the dermis. Triamcinolone intralesional injection, oral methylprednisolone and doxycycline was not effective. Seven months after the first visit, stereotactic radiosurgery was performed for cavernous hemangioma. We prescribed the same dose of methylprednisolone as before. Despite dosage was not increased, symptoms tended to improve. After 2 months, oral methylprednisolone was discontinued and intermittent triamcinolone intralesional injection was performed solely. Her skin condition is improving continuously.





